arXiv:1412.0302v1 [math.QA] 30 Nov 2014

INTEGRABLE CLUSTERS

ARKADY BERENSTEIN, JACOB GREENSTEIN, AND DAVID KAZHDAN

Abstract. The goal of this note is to study quantum clusters in which cluster variables (not coefficients) commute which each other. It turns out that this property is preserved by mutations. Remarkably, this is equivalent to the celebrated sign coherence conjecture recently proved by M. Gross, P. Hacking, S. Keel and M. Kontsevich in [6].

1. Main results

Let $\tilde{B}$ be an integer $m \times n$ matrix, $n \leq m$, and let $\Lambda = (\lambda_{ij})_{1 \leq i,j \leq m}$ be a rational skew-symmetric $m \times m$-matrix compatible with $\tilde{B}$ in the sense of [3], that is

$$\tilde{B}^T \Lambda = \begin{pmatrix} D & \mathbf{0} \end{pmatrix} \tag{1.1}$$

where D is a rational invertible diagonal $n \times n$-matrix. This defines, on the one hand, a seed $\Sigma = (\mathbf{x}, \tilde{B})$, $\mathbf{x} = (x_1, \dots, x_m)$ and the upper cluster algebra $\mathcal{U} = \mathcal{U}(\Sigma) \subset \mathscr{L}_m = \mathbb{Q}[x_1^{\pm 1}, \dots, x_m^{\pm 1}]$ and, on the other hand, the Poisson algebra structure on $\mathscr{L}_m$ via $\{x_i, x_j\}_\Lambda = \lambda_{ij} x_i x_j$, $1 \leq i, j \leq m$ so that $\mathcal{U}$ is a Poisson subalgebra of $\mathscr{L}_m$. This in turn defines a Poisson scheme $\mathscr{X}$ such that for any field extension $\mathbb{F}$ of $\mathbb{Q}$, $\mathscr{X}_{\mathbb{F}} = \mathrm{Spec}(\mathcal{U} \otimes_{\mathbb{Q}} \mathbb{F})$.

We say that the seed $\Sigma = (\mathbf{x}, \tilde{B})$ is *integrable* if $\{x_i, x_j\}_\Lambda = 0$ for all $1 \leq i, j \leq n$ for some Λ compatible with $\tilde{B}$. It is easy to show (Lemma 2.4) that if Σ is principal (see Section 2) then it is integrable and the matrix Λ is uniquely determined by $\tilde{B}$ and D. We say that Σ is *completely integrable* if Σ and all its mutations are integrable. This definition is justified by the following observation. Let $m = 2n$ and assume that Σ is integrable and that Λ is invertible, i.e., $\{\cdot, \cdot\}_\Lambda$ is a symplectic bracket. Then for any $H \in \mathbb{Q}[x_1, \dots, x_n] \setminus \{0\}$ the triple $(\mathcal{U}, \{\cdot, \cdot\}_\Lambda, H)$ is an integrable system with Hamiltonian H; the natural map $\pi_\Sigma : \mathscr{X} \to \mathbb{A}^n$ is a Lagrangian fibration (see [1]).

Theorem 1.1. *Every principal seed is completely integrable.*

Corollary 1.2. *For any seed $\Sigma' = (\mathbf{y}, \tilde{B}')$, $\mathbf{y} = (y_1, \dots, y_m)$, mutation equivalent to a principal integrable seed Σ, the natural inclusion $\Bbbk[y_1, \dots, y_n] \subset \mathcal{U}(\Sigma') = \mathcal{U}$ defines a Lagrangian fibration $\pi_{\Sigma'} : \mathscr{X} \to \mathbb{A}^n$.*

This gives rise to the following natural problem.

Problem 1.3. *Given mutation equivalent integrable seeds Σ, Σ', describe the intersection $\pi_\Sigma^{-1}(c) \cap \pi_{\Sigma'}^{-1}(c')$ for generic points $c, c' \in \mathbb{A}^n$ and determine for which Σ, Σ' this intersection is transversal.*

It turns out that classical results carry over verbatim to the quantum case. We say that a quantum seed $\Sigma = (\mathbf{X}, \tilde{B})$, where $\mathbf{X} = (X_1, \dots, X_m)$ and $\tilde{B}$ is an $m \times n$ integer matrix (see Section 2), is *integrable* if $X_i X_j = X_j X_i$ for all $1 \leq i, j \leq n$. Then, similarly to the classical case, we say that a quantum seed Σ is *completely integrable* if Σ and all its mutations are integrable.

Theorem 1.4. *Every integrable principal quantum seed is completely integrable.*

By Lemma 2.6 any (quantum or classical) seed can be converted into a principal integrable one merely by duplicating the ambient (quantum) torus as in [3, Section 3].

A *generalized quantum integrable system* is a pair $(\mathcal{U}, \iota)$ where $\mathcal{U}$ is a $\Bbbk$-algebra of Gelfand-Kirillov dimension $2n$ and $\iota : \Bbbk[x_1, \dots, x_n] \hookrightarrow \mathcal{U}$ is an embedding of algebras such that $\iota(\Bbbk[x_1, \dots, x_n])$ is a maximal commutative subalgebra of $\mathcal{U}$. This defines a quantum analogue of Lagrangian fibration, whose *quantum Lagrange fiber* over a maximal ideal $\mathfrak{m}$ of $\Bbbk[x_1, \dots, x_n]$ is the left $\mathcal{U}$-module $\mathcal{U}_{\iota,\mathfrak{m}} := \mathcal{U}/\mathcal{U} \cdot \iota(\mathfrak{m})$. Then the (quantum) intersection of $\mathcal{U}_{\iota,\mathfrak{m}}$ and $\mathcal{U}_{\iota',\mathfrak{m}'}$ is the left $\mathcal{U}$-module $\mathcal{U}/\mathcal{U} \cdot (\iota(\mathfrak{m}) + \iota'(\mathfrak{m}'))$.

Following [2], given a field $\Bbbk$ containing $\mathbb{Q}(q^{\frac{1}{2}})$ and a *quantum seed* $\Sigma = (\mathbf{X}, \tilde{B})$, we denote by $\mathcal{U} = \mathcal{U}(\Sigma)$ its *upper quantum cluster algebra* (see Section 2 for the details).

Corollary 1.5. *For any quantum seed $\Sigma' = (\mathbf{Y}, \tilde{B}')$ mutation equivalent to a given principal integrable quantum seed Σ, the natural inclusion $\iota : \Bbbk[Y_1, \dots, Y_n] \hookrightarrow \mathcal{U}(\Sigma') = \mathcal{U}$ defines a generalized quantum integrable system $(\mathcal{U}, \iota)$.*

Date: December 2, 2014.

The authors were partially supported by the BSF grant no. 2012365 (A. B. and D. K.), NSF grant DMS-1403527 (A. B.), the ERC grant no. 247049 (D.K.) and the Simons foundation collaboration grant no. 245735 (J. G.) .

2. Notation and proofs

We will only prove quantum results since their classical counterparts follow by specializing q to 1.

Let $\Lambda \in \mathrm{Mat}_{m\times m}(\mathbb{Z})$ be compatible with $\tilde{B}$ in the sense of (1.1) where $D \in \mathrm{Mat}_{n\times n}(\mathbb{Z})$ and has positive diagonal entries. Following [2], we associate with the pair $(\Lambda, \tilde{B})$ a *quantum seed* $\Sigma = (\mathbf{X}, \tilde{B})$, $\mathbf{X} = (X_1, \dots, X_m) \in (\mathcal{F}^\times)^m$ where $\mathcal{F}$ is a skew field, such that the subalgebra $\mathscr{L}_{\mathbf{X}}$ of $\mathcal{F}$ generated by $\mathbf{X}$ over some central subfield $\Bbbk$ of $\mathcal{F}$ containing $\mathbb{Q}(q^{\frac{1}{2}})$ has presentation

$$X_iX_j = q^{\lambda_{ij}}X_jX_i, \qquad 1 \le i < j \le m.$$

Given $1 \le j \le n$, we define $\mu_j(\Sigma) = (\mathbf{X}', \mu_j(\tilde{B}))$ where $\mu_j(\tilde{B})$ is the Fomin-Zelevinsky mutation of $\tilde{B}$ from [4] and $\mathbf{X}'$ is obtained from $\mathbf{X}$ by replacing X_j with

$$X_j' = X^{[b_j]_+ - e_j} + X^{[-b_j]_+ - e_j}, \tag{2.1}$$

where b_j is the jth column of $\tilde{B}$, for each $a = (a_1, \dots, a_m) \in \mathbb{Z}^m$ we set $[a]_+ = (\max(0, a_1), \dots, \max(0, a_m))$,

$$X^a = q^{\frac{1}{2}\sum_{1\le i<j\le m}\lambda_{ji}a_ia_j}X_1^{a_1}\cdots X_m^{a_m}$$

and $\{e_i\}_{1\le i\le m}$ is the standard basis of $\mathbb{Z}^m$. After [2, Section 2], $\mu_j(\Sigma)$ is also a quantum seed and we refer to it as the jth mutation of Σ. A quantum seed Σ' is *mutation equivalent* to Σ if it can be obtained from Σ by a sequence of mutations.

Following [7], we say that an $m \times n$-matrix $\tilde{B}$ is *sign-coherent* if for every $1 \le j \le n$ there exists $\epsilon_j \in \{-1, 1\}$ such that $\epsilon_j b_{ij} \ge 0$ for all $n+1 \le i \le m$. We say that $\tilde{B}$ is *totally sign-coherent* if all matrices mutation equivalent to $\tilde{B}$ are sign-coherent.

Proposition 2.1. *Suppose $\Lambda \in \mathrm{Mat}_{m\times m}(\mathbb{Z})$ and $\tilde{B} \in \mathrm{Mat}_{m\times n}(\mathbb{Z})$ are compatible, $\tilde{B}$ is totally sign-coherent and the corresponding quantum seed Σ is integrable. Then Σ is completely integrable.*

Proof. We need the following

Lemma 2.2. *Suppose that $\Sigma = (\mathbf{X}, \tilde{B})$ is integrable and $\tilde{B}$ is sign-coherent. Then $\mu_j(\Sigma)$, $1 \le j \le n$, is integrable.*

Proof. Since $\tilde{B}$ is sign-coherent, either $X_jX^{[b_j]_+-e_j}$ or $X_jX^{[-b_j]_+-e_j}$ is contained in $\Bbbk[X_1, \dots, X_n]$. Since for every $1 \le i \ne j \le n$ we have $X_iX^{[b_j]_+-e_j} = q_{ij}X^{[b_j]_+-e_j}X_i$ and $X_iX^{[-b_j]_+-e_j} = q_{ij}X^{[-b_j]_+-e_j}X_i$ for some $q_{ij} \in \Bbbk^\times$, it follows that $q_{ij} = 1$. Then by (2.1) we have $X_iX_j' = X_j'X_i$ for all $1 \le i \ne j \le n$. □

We complete the proof by induction on the number of mutations applied to the initial seed Σ. If $\Sigma' = \mu_{j_1}\cdots\mu_{j_k}(\Sigma) = \mu_{j_1}(\Sigma'')$ where $\Sigma'' = (\mathbf{X}'', \tilde{B}'') = \mu_{j_2}\cdots\mu_{j_k}(\Sigma)$ is integrable by the induction hypothesis and $\tilde{B}''$ is sign-coherent by assumption on $\tilde{B}$. It remains to apply the Lemma. □

Proof of Theorem 1.4. Recall from [5, Remark 3.2] that $\tilde{B}$ is called principal if $\tilde{B} = \begin{pmatrix} B \\ I_n \end{pmatrix}$ where $B \in \mathrm{Mat}_{n\times n}(\mathbb{Z})$ and DB is skew symmetric for some $D \in \mathrm{Mat}_{n\times n}(\mathbb{Z})$ diagonal with positive diagonal entries. The following result was initially conjectured in [5, Conjecture 5.4 and Proposition 5.6(iii)] (see also [7, (1.8)]).

Lemma 2.3 ([6, Corollary 5.5])**.** *Any principal $\tilde{B}$ is totally sign-coherent.*

Theorem 1.4 is immediate from Proposition 2.1 and the above Lemma. □

Proof of Corollary 1.5. We need the following obvious classification of (quantum) integrable seeds with $m = 2n$.

Lemma 2.4. *Let*

$$\Lambda = \begin{pmatrix} \mathbf{0} & \Lambda_1 \\ -\Lambda_1^T & \Lambda_2 \end{pmatrix}, \qquad \tilde{B} = \begin{pmatrix} B \\ C \end{pmatrix} \tag{2.2}$$

with $B, C, \Lambda_1, \Lambda_2 \in \mathrm{Mat}_{n\times n}(\mathbb{Q})$ and $\Lambda_2^T = -\Lambda_2$. Then (1.1) holds for some invertible diagonal $D \in \mathrm{Mat}_{n\times n}(\mathbb{Q})$ if and only if

$$\det C \ne 0, \quad (DB)^T = -DB, \quad \Lambda_1 = -DC^{-1}, \quad \Lambda_2 = -(C^{-1})^TDBC^{-1}.$$

Furthermore, following [2], to each quantum seed Σ one associates the *quantum upper cluster algebra* $\mathcal{U}(\Sigma) = \bigcap_{1\le i\le n}\mathcal{U}_i$ where $\mathcal{U}_i$ is the subalgebra of $\mathscr{L}_{\mathbf{X}}$ generated by $\mathbf{X}$ and X_i'. As shown in [2, Theorem 5.1], $\mathcal{U}(\Sigma) = \mathcal{U}(\mu_j(\Sigma))$ for all $1 \le j \le n$. We need the following Lemma.

Lemma 2.5. *Let $\Sigma = (\mathbf{X}, \tilde{B})$ be a quantum integrable seed with $m = 2n$. Then*

(a) *$\Bbbk[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$ is a maximal commutative subalgebra of $\mathscr{L}_{\mathbf{X}}$;*

(b) $\Bbbk[X_1, \dots, X_n]$ *is a maximal commutative subalgebra of* $\mathcal{U}(\Sigma)$.

Proof. To prove (a) note that for each $1 \le i \le n$ the centralizer $\mathscr{C}_i$ of X_i in $\mathscr{L}_{\mathbf{X}}$ is the $\Bbbk$-linear span of $\{X^a : \sum_{1\le j\le m} \lambda_{ij} a_j = 0\}$. Since Λ is as in (2.2) with $\det \Lambda_1 \neq 0$, $\bigcap_{1\le i\le n} \mathscr{C}_i$ is spanned by all monomials X^a with $a_j = 0$, $j > n$. This implies that $\Bbbk[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$ is a maximal commutative subalgebra of $\mathscr{L}_{\mathbf{X}}$.

To prove (b), denote $\mathcal{U}_i' = \mathcal{U}_i \cap \Bbbk[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$. It is easy to see that $\mathcal{U}_i' = \Bbbk[X_1^{\pm 1}, \dots, X_i, \dots, X_n^{\pm 1}]$. Since $\mathcal{U} \cap \Bbbk[X_1^{\pm 1}, \dots, X_n^{\pm 1}] = \bigcap_{1\le i\le n} \mathcal{U}_i' = \Bbbk[X_1, \dots, X_n]$, part (b) is now immediate from part (a). □

The Corollary 1.5 is now immediate because each quantum seed mutation equivalent to a given principal integrable one is automatically integrable by Theorem 1.4. □

We conclude by showing that every (quantum) seed can be converted into a principal completely integrable one. Recall that in [3, Section 3] to every seed $\Sigma = (\mathbf{X}, \tilde{B})$ in $\mathscr{L}_{\mathbf{X}}$ one associates a seed $\Sigma^\bullet = (\mathbf{X}^\bullet, \tilde{B}^\bullet)$ in the duplicated quantum torus $\mathscr{L}_{\mathbf{X}}^{(2)} = \bigoplus_{e,e'\in\mathbb{Z}^m} \Bbbk X^{(e,e')}$ with $\Lambda^{(2)} = \begin{pmatrix} \Lambda & \mathbf{0} \\ \mathbf{0} & -\Lambda \end{pmatrix}$, as follows

$$X_i^\bullet = X^{(e_i, e_i)}, \qquad X_{i+n}^\bullet = X^{(b_i^{>n}, -b_i^{\le n})}, \qquad 1 \le i \le n, \qquad \tilde{B}^\bullet = \begin{pmatrix} B \\ I_n \end{pmatrix}$$

where we abbreviate $a^{>n} = \sum_{n<i\le m} a e_i$, $a^{\le n} = \sum_{1\le i\le n} a_i e_i$ for $a = (a_1, \dots, a_m) \in \mathbb{Z}^m$. By definition, $\mathscr{L}_{\mathbf{X}}$ identifies with a subalgebra of $\mathscr{L}_{\mathbf{X}}^{(2)}$ via $X^e \mapsto X^{(e,0)}$, $e \in \mathbb{Z}^m$. Then $\mathscr{L}_{\mathbf{X}}^{(2)} = \mathscr{L}_{\mathbf{X}} \cdot \mathscr{C}_{\mathbf{X}}$ and hence $\mathcal{U}(\Sigma) \cdot \mathscr{C}_{\mathbf{X}} = \mathcal{U}(\Sigma^\bullet) \cdot \mathscr{C}_{\mathbf{X}}$ as subalgebras of $\mathscr{L}_{\mathbf{X}}^{(2)}$, where $\mathscr{C}_{\mathbf{X}}$ is the subalgebra of $\mathscr{L}_{\mathbf{X}}^{(2)}$ generated by the $X^{(e_i,0)}$, $n < i \le m$ and by the $X^{(0,e_j)}$, $1 \le j \le m$.

The following is immediate from [3, (3.19)–(3.22) and Lemma 3.4].

Lemma 2.6. *The seed* $\Sigma^\bullet$ *is principal and integrable, hence completely integrable.*

References

[1] V. I. Arnol′d, *Mathematical methods of classical mechanics*, Graduate Texts in Mathematics, vol. 60, Springer-Verlag, 1989.
[2] A. Berenstein and A. Zelevinsky, *Quantum cluster algebras*, Adv. Math. **195** (2005), no. 2, 405–455.
[3] ———, *Triangular bases in quantum cluster algebras*, Int. Math. Res. Not. **2014** (2014), no. 6, 1651–1688.
[4] S. Fomin and A. Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529.
[5] ———, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), no. 1, 112–164.
[6] M. Gross, P. Hacking, S. Keel, and M. Kontsevich, *Canonical bases for cluster algebras*, available at `arXiv:1411.1394`.
[7] T. Nakanishi and A. Zelevinsky, *On tropical dualities in cluster algebras*, Algebraic groups and quantum groups, Contemp. Math., vol. 565, Amer. Math. Soc., Providence, RI, 2012, pp. 217–226.

Department of Mathematics, University of Oregon, Eugene, OR 97403, USA
E-mail address: `arkadiy@math.uoregon.edu`

Department of Mathematics, University of California, Riverside, CA 92521.
E-mail address: `jacob.greenstein@ucr.edu`

Department of Mathematics, Heberew University, Jerusalem, Israel
E-mail address: `kazhdan@math.huji.ac.il`